

March 1993

2D dilaton-Maxwell gravity as a fixed point of the renormalization group

E. ELIZALDE¹

*Department E.C.M., Faculty of Physics, University of Barcelona,
Diagonal 647, 08028 Barcelona, Spain*

S. NAFTULIN

Institute for Single Crystals, 60 Lenin Ave., 310141 Kharkov, Ukraine

and

S.D. ODINTSOV²

*Tomsk Pedagogical Institute, 634041 Tomsk, Russia,
and Department of Physics, Faculty of Science, Hiroshima University,
Higashi-Hiroshima 724, Japan*

Abstract

A general model of dilaton-Maxwell gravity in two dimensions is investigated. The corresponding one-loop effective action and the generalized β -functions are obtained. A set of models that are fixed points of the renormalization group equations are presented.

¹E-mail address: eli @ ebubecm1.bitnet

² E-mail address: odintsov @ theo.phys.sci.hiroshima-u.ac.jp

Recently, the issue of the Hawking evaporation of black holes [1] has been discussed in two-dimensional (2D) dilaton gravity with matter, what is considered to be a toy model for 2D gravity. Based on ref. [2], different aspects for different models of 2D quantum dilaton gravity—including relevant questions about the associated black holes—have been extensively considered in the recent literature (for a review and a quite comprehensive list of references see, for example, [3]). Due to the wide variety of models of 2D gravity under current investigation (some of which are classically equivalent) the following question arises naturally: what is a good criterion in order to select reasonable models of 2D gravity?

In the present letter we suggest one: to use the renormalization group as such a criterion—in a sense that will become clear in the progress of this work. We start from the general model of 2D gravity defined by the action

$$S = - \int d^2x \sqrt{g} \left[\frac{1}{2} Z(\Phi) g^{\mu\nu} \partial_\mu \Phi \partial_\nu \Phi + C(\Phi) R + V(\Phi) + \frac{1}{4} f(\Phi) F_{\mu\nu}^2 \right], \quad (1)$$

where Φ is the dilaton field, $Z(\Phi)$, $C(\Phi)$, $V(\Phi)$ and $f(\Phi)$ are arbitrary functions, and $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$. Notice that the model (1) is related, via some compactification, with the four-dimensional Einstein-Maxwell theory, which admits charged black hole solutions [4]. For some specific values of the functions in (1), this general model corresponds, in fact, to the heterotic string effective action [5,6]. Notice also that the theory (1) is multiplicatively renormalizable [8,9].

The one-loop effective action for theory (1) with the choice: $Z(\Phi) = 1$, $C(\Phi) = \text{const.}$, and $V(\Phi)$ and $f(\Phi)$ arbitrary, has been calculated in different gauges in ref. [7]. Here, we are going to find the one-loop effective action using the background field formalism for this model (1), in the general case.

In the background field method, we split

$$g_{\mu\nu} \longrightarrow \bar{g}_{\mu\nu} = g_{\mu\nu} + h_{\mu\nu}, \quad \Phi \longrightarrow \bar{\Phi} = \Phi + \varphi, \quad A_\mu \longrightarrow \bar{A}_\mu = A_\mu + Q_\mu, \quad (2)$$

where $h_{\mu\nu}$, φ and Q_μ are the quantum fields. The simplest minimal gauge fixing action is given by

$$S_{GF} = -\frac{1}{2} \int d^2x \chi^A c_{AB} \chi^B, \quad (3)$$

where $A \equiv \{\mu, *\}$, $c_{\mu\nu} = -C \sqrt{g} g_{\mu\nu}$, $\chi^\mu = -\nabla_\nu \bar{h}^{\mu\nu} + \frac{C'}{C} \nabla^\mu \varphi$, $C_* = \sqrt{g} f$, $\chi_* = -\nabla^\nu Q_\nu$. Working in this gauge we obtain the following effective action (details of the very involved

calculations will be reported elsewhere):

$$\begin{aligned}\Gamma_{div} = & -\frac{1}{2\epsilon} \int d^2x \sqrt{g} \left\{ 5R + \frac{2}{C}V + \frac{2}{C'}V' + \left(\frac{f'}{2C'} - \frac{f}{2C} \right) F^2 \right. \\ & + \left(\frac{f'}{f} + \frac{C'}{C} - \frac{Z}{C'} \right) (\Delta\Phi) \\ & \left. + \left(\frac{f''}{f} - \frac{f'^2}{f^2} + \frac{C''}{C} - 3\frac{C'^2}{C^2} - \frac{C''Z}{C'^2} \right) (\nabla^\lambda\Phi)(\nabla_\lambda\Phi) \right\}. \quad (4)\end{aligned}$$

It must be remarked that in this expression all surface divergent terms have been kept.

Let us discuss the on-shell limit of the effective action (4). The corresponding classical field equations are

$$\begin{aligned}\frac{\delta S}{\delta\Phi} &= -\nabla_\nu(Zg^{\mu\nu}\partial_\mu\Phi) + \frac{1}{2}Z'(\nabla^\mu\Phi)(\nabla_\mu\Phi) + C'R + V' + \frac{1}{4}f'F^2 = 0, \\ g^{\mu\nu}\frac{\delta S}{\delta g^{\mu\nu}} &= -\Delta C + V - \frac{1}{4}fF^2 = 0,\end{aligned} \quad (5)$$

and substituting eqs. (5) into the effective action (4), keeping all the surface counterterms, we obtain

$$\Gamma_{div}^{on-shell} = -\frac{1}{2\epsilon} \int d^2x \sqrt{g} \left\{ 3R + \Delta [\ln(fC^3)] + \nabla^\lambda \left[\frac{Z}{C'} \nabla_\lambda(\Phi) \right] \right\}. \quad (6)$$

As we see, the theory considered is finite on-shell.

We will now study the renormalization structure and renormalization group equations for dilaton gravity with the action (1). One could, of course, discuss the renormalization of the metric and find the restrictions imposed by off-shell multiplicative renormalizability (in the usual sense) on the form of the functions $Z(\Phi)$, $C(\Phi)$, $V(\Phi)$ and $f(\Phi)$. However, instead of doing this, we will not renormalize the fields but rather will consider the functions Z , C , V and f as generalized effective couplings (generalized renormalizability). Then the generalized β -functions can be found and generalized renormalization group equations are generated.

The general structure of renormalization for general couplings is given by

$$Z_0 = (\mu^2)^{\epsilon'} \left[Z + \sum_{k=1}^{\infty} \frac{a_{kZ}(Z, C, V, f)}{\epsilon'^k} \right], \quad (7)$$

where $\epsilon' = n - 2$, and similar expressions for C , V and f . As it follows from one-loop renormalization, eq. (4),

$$a_{1Z} = -\frac{Z'}{C'} + \frac{2C''^2}{C^2} + \frac{2C''Z}{C'^2}, \quad a_{1C} = 0, \quad a_{1V} = -\frac{V}{C} - \frac{V'}{C'}, \quad a_{1f} = \frac{f}{C} - \frac{f'}{C'}. \quad (8)$$

Now, the generalized one-loop β -functions are given by the standard relations:

$$\beta_T = -a_{1T} + Z \frac{\delta a_{1T}}{\delta Z} + C \frac{\delta a_{1T}}{\delta C} + V \frac{\delta a_{1T}}{\delta V} + f \frac{\delta a_{1T}}{\delta f}, \quad (9)$$

where $T \equiv \{Z, C, V, f\}$. Using (8) and (9) we obtain

$$\begin{aligned} \beta_C &= 0, \\ \beta_V &= \frac{V}{C} + \frac{V'}{C'} - \frac{VC''}{C'^2} - \frac{CV''}{C'^2} + 2 \frac{CV'C''}{C'^3} \\ \beta_Z &= \frac{Z'}{C'} + \frac{2C'^2}{C^2} - \frac{4C''}{C} - \frac{ZC''}{C'^2} + 3 \frac{CZ''}{C'^2} - 2 \frac{CZ'C''}{C'^3}, \\ \beta_f &= -\frac{f}{C} + \frac{f'}{C'} - \frac{fC''}{C'^2} - \frac{Cf''}{C'^2} + 2 \frac{Cf'C''}{C'^3}. \end{aligned} \quad (10)$$

The renormalization group equations have the following form:

$$\frac{\partial C}{\partial t} = \beta_C, \quad \frac{\partial V}{\partial t} = \beta_V, \quad \frac{\partial Z}{\partial t} = \beta_Z, \quad \frac{\partial f}{\partial t} = \beta_f, \quad (11)$$

with $t = \ln \mu^2$. The system of equations (11) is very difficult to solve. It depends not only on the scaling parameter t , as in usual field theory, but also on some unknown functions of the field variables. Moreover, nobody has any idea about the proper boundary conditions (initial data) that the partial differential equation system (11) should satisfy.

Notwithstanding that, we can extract some useful information yet from the renormalization group equations (11). In particular, we can look for fixed points of this system, what does not at all involve the knowledge of initial data. The equations they must satisfy are

$$\beta_C = 0, \quad \beta_V = 0, \quad \beta_Z = 0, \quad \beta_f = 0. \quad (12)$$

The system of differential equations (12) is still very complicated. Nevertheless, some basic, particular solutions of the same can be discovered. For example, motivated by the CGHS action [2], we can look for fixed points of the following type:

$$C = e^{a_1 \Phi}, \quad V = e^{a_2 \Phi}, \quad Z = e^{a_3 \Phi}, \quad f = e^{a_4 \Phi}, \quad (13)$$

where a_1, a_2, a_3 and a_4 are some constants. Substituting (13) into eq. (12) we obtain the following equations:

$$\begin{aligned} a_1 &= \frac{a_2(a_2 - 1)}{a_2 + 1}, \\ 3a_3^3 + a_3(1 - 3a_1) + 2a_1(1 - 2a_1) &= 0, \\ a_4^2 - 3a_1a_4 + 2a_1^2 &= 0, \end{aligned} \quad (14)$$

which have the solution

$$\begin{aligned} a_1 = 0, \quad a_2 = 1, \quad a_3 = 0, -\frac{1}{3}, \quad a_4 = 0; \\ a_1 = \frac{2}{3}, \quad a_2 = 2, \quad a_3 = \frac{1}{6} \left(1 \pm \sqrt{19/3} \right), \quad a_4 = \frac{2}{3}, \frac{4}{3}; \quad \text{etc.} \end{aligned} \quad (15)$$

In the same way, different particular cases of fixed points can be considered. For instance, for

$$C = \Phi^{\alpha_1}, \quad V = \Phi^{\alpha_2}, \quad Z = \Phi^{\alpha_3}, \quad f = \Phi^{\alpha_4}, \quad (16)$$

we get

$$\begin{aligned} \alpha_1(\alpha_1 - 2) &= 0, \\ \alpha_2^2 + (1 - 3\alpha_1)\alpha_2 - \alpha_1 &= 0, \\ 3\alpha_3^2 - (1 + \alpha_1)\alpha_3 - \alpha_1(\alpha_1 - 1) &= 0, \\ \alpha_4^2 + (1 - 3\alpha_1)\alpha_4 + \alpha_1(2\alpha_1 - 1) &= 0. \end{aligned} \quad (17)$$

Particular solutions are

$$\begin{aligned} \alpha_1 = 0, \quad \alpha_2 = -1, \quad \alpha_3 = \frac{1}{3}, \quad \alpha_4 = -1; \\ \alpha_1 = 2, \quad \alpha_2 = \frac{1}{2} \left(5 \pm \sqrt{33} \right), \quad \alpha_3 = \frac{1}{2} \left(5 \pm \sqrt{11/3} \right), \quad \alpha_4 = 2, 3; \quad \text{etc.} \end{aligned} \quad (18)$$

Guided by the CGHS model, one could also try to find 'mixed' solutions of the type

$$C = \Phi^{a_1} e^{b_1 \Phi}, \quad V = \Phi^{a_2} e^{b_2 \Phi}, \quad Z = \Phi^{a_3} e^{b_3 \Phi}, \quad f = \Phi^{a_4} e^{b_4 \Phi}. \quad (19)$$

It can be shown, however, that no new solution is found in this way: the only possibilities reduce to the cases above (namely, either all a_i or all b_i vanish).

Thus, we have shown that (at least in principle) one can find ultraviolet stable fixed points for the generalized couplings $Z(\Phi)$, $C(\Phi)$, $V(\Phi)$ and $f(\Phi)$.

Summing up, we have here investigated the one-loop renormalization structure of a general model of 2D dilaton-Maxwell gravity (1), and the divergences of the corresponding one-loop effective action have been found. Generalized renormalizability of the general model of dilaton gravity has been discussed, and the corresponding generalized β functions have been obtained.

The analysis of the renormalization group equations yields some set of generalized couplings $\{Z, C, V, f\}$ which are fixed points of such equations. It is interesting to notice

that the CGHS model does *not* belong to this set. A further remark is the fact that if one requires the renormalizability of the theory in the usual sense, then one can renormalize the metric of the theory (1) through the following transformation

$$g_{\mu\nu} = \exp \left[\frac{1}{\epsilon} \left(\frac{1}{C} + \frac{Z}{2C'^2} \right) \right] g_{\mu\nu}^R. \quad (20)$$

It turns out that the 2D dilaton-Maxwell theory (1) is multiplicatively renormalizable off-shell in the usual sense for the following choice of potentials:

$$V = \exp \left[aC + \int \frac{Z d\Phi}{2C'} \right], \quad f = \exp \left[-bC - \int \frac{Z d\Phi}{2C'} \right], \quad (21)$$

where a and b are arbitrary constants. For the theory of ref. [7] the above potentials are of Liouville type, what is in full agreement with ref. [7]. We observe, finally, that the condition of off-shell renormalizability in the usual sense can be considered also as a good criterion for selecting 2D gravity models.

Acknowledgments

We are grateful to I. Antoniadis, F. Englert, Y. Kazama, H. Osborn, A.A. Slavnov and I.V. Tyutin for useful discussions at different stages of this work. S.D.O. wishes to thank the Japan Society for the Promotion of Science (JSPS, Japan) for financial support and the Particle Physics Group at Hiroshima University for kind hospitality. E.E. has been supported by DGICYT (Spain), research project PB90-0022, and by the Generalitat de Catalunya.

References

- [1] S.W. Hawking, Commun. Math. Phys. **43** (1975) 199.
- [2] C.G. Callan, S.B. Giddings, J.A. Harvey and A. Strominger, Phys. Rev. **D45** (1992) 1005.
- [3] J.A. Harvey and A. Strominger, preprint EFI-92-41 (1992).
- [4] G.W. Gibbons, Nucl. Phys. **B207** (1982) 337; G.W. Gibbons and K. Maeda, Nucl. Phys. **B298** (1988) 74.
- [5] C.G. Callan, D. Friedan, E. Martinec and M.J. Perry, Nucl. Phys. **B262** (1985) 593; C.G. Callan, I. Klebanov and M.J. Perry, Nucl. Phys. **B278** (1986) 78.
- [6] M. McGuigan, C. Nappi and S. Yost, IAS preprint IASSNS- HEP-91157 (1991).
- [7] E. Elizalde and S.D. Odintsov, Mod. Phys. Lett. **A8** (1993) 33; E. Elizalde, S. Naf-tulin and S.D. Odintsov, preprint HUPD-93-03 (1993).
- [8] T. Banks and M. O'Loughlin, Nucl. Phys. **B362** (1991) 649.
- [9] S.D. Odintsov and I.L. Shapiro, Phys. Lett. **B263** (1991) 183.